

Conjugacy problem for subgroups with applications to Artin groups and braid type groups

NUNO FRANCO

ABSTRACT. Let G be a group endowed with a solution to the conjugacy problem and with an algorithm which computes the centralizer in G of any element of G . Let H be a subgroup of G . We give some conditions on H , under which we provide a solution to the conjugacy problem in H . We apply our results to some Artin groups and braid type groups. In particular, we give explicit solutions to the conjugacy problem in the Artin groups of type $\tilde{A}_n$ and $\tilde{C}_n$.

February 1, 2008

AMS Subject classification: Primary 20F36, Secondary 20F10.

1. INTRODUCTION

Let G be a group. A *solution for the conjugacy problem* in G consists on an algorithm which, for given $a, b \in G$, determines whether there exists $c \in G$ such that $a = c^{-1}bc$. An *explicit solution for the conjugacy problem* is a solution for the conjugacy problem that also determines an element $c \in G$ such that $a = c^{-1}bc$. Define the *centralizer*, $C_G(a) = C(a)$, of an element $a \in G$ as the set of elements in G that commute with a . A solution for the *membership problem* in G is an algorithm that, for a given finitely generated subgroup $H < G$ and a given $g \in G$, decides whether $g \in H$.

Throughout this paper G will denote a group endowed with an explicit solution for the conjugacy problem and with an algorithm that, for any element $a \in G$, computes a generating set for the centralizer of a in G (which we always assume to be finitely generated).

The present paper concerns the following question: under which conditions, for a given $H < G$, we can find an explicit solution to the conjugacy problem in H .

Let $H < G$ be a subgroup. We say that H verifies Condition **PC** if there exists a group K , a subgroup K' of K and an homomorphism $\phi : G \longrightarrow K$, such that:

- $H = \phi^{-1}(K')$,
- there is a solution for the membership problem in K , and
- the subgroup K' is finite.

Let H be a subgroup of G that verifies Condition **PC**. Our goal is to compute an explicit solution to the conjugacy problem in H , using the triple (K, K', ϕ) of the definition.

Remark that Condition **PC** is essentially a condition on the group K and not on the subgroup H . This condition on K is very restrictive since there are few known classes of groups in which we can give a solution to the membership problem. It is known that the following classes of groups have a solution to the membership problem:

- finite groups,
- free groups,
- abelian groups,
- recursively presented metabelian groups,
- finitely generated fully residually free groups.

The existence of a solution to the conjugacy problem is also closely related to the group based cryptographic systems. In [2] and [15] are introduced some cryptographic systems, called group based cryptographic system, in which the encryption security is based on:

- the complexity of a solution to the conjugacy problem, and
- the existence of a fast solution to the word problem.

A subgroup H for which it does not exists a triple that verifies Condition **PC** will assure us that it is not possible to use our approach to attack an H -based cryptographic system. Even if we use a subgroup H of G that verifies Condition **PC** to implement a cryptographic system, we still have, in general, some security advantages when comparing to an G -based cryptographic system:

- the same fast solution for the word problem in H as in the group G , and
- a higher complexity of the solution of the conjugacy problem in H .

In Section 2 we present some examples of groups and subgroups in which we can apply or not our theory. Sections 3 contains the main results and algorithms regarding the conjugacy problem. Finally, in Section 4, we make some brief remarks about the complexity of the algorithms presented in this paper.

2. APPLICATIONS

In this section we will give some examples of groups in which we can apply our theory. These examples will be subgroups of Artin groups, and other groups that admit braid pictures. We start with some definitions.

Define a *Coxeter matrix* of rank n as a $n \times n$ symmetric matrix, $M = (m_{i,j})$, that verifies: $m_{i,i} = 1$ for all $i = 1, \dots, n$, and $m_{i,j} \in \{2, 3, \dots\} \cup \{\infty\}$, for all $i, j \in \{1, \dots, n\}$, $i \neq j$. Let M be a Coxeter matrix. The *Coxeter graph* associated to M is the labelled graph, Γ , defined as follows. The set of vertices of Γ is $\{1, \dots, n\}$. If $m_{i,j} = 2$ then there is no edge between i and

j , if $m_{i,j} = 3$, then there is a non-labelled edge between i and j , and, finally, if $m_{i,j} > 3$ or $m_{i,j} = \infty$, then there is an edge between i and j labelled by $m_{i,j}$.

Let G be a group. For $a, b \in G$ and $n \in \mathbb{N}$ we define the word

$$\text{prod}_n(a, b) \begin{cases} (ab)^{\frac{n}{2}} & \text{if } n \equiv 0 \pmod{2} \\ (ab)^{\frac{n-1}{2}}a & \text{if } n \equiv 1 \pmod{2} \end{cases}$$

Define the *Coxeter group* W associated to the Coxeter graph Γ as the group presented by

$$W = W(\Gamma) = \left\langle s_1, s_2, \dots, s_n \mid \begin{array}{l} s_i^2 = 1, \quad i \in \{1, \dots, n\} \\ \text{prod}_{m_{i,j}}(s_i, s_j) = \text{prod}_{m_{j,i}}(s_j, s_i) \\ \quad i \neq j \text{ and } m_{i,j} \neq \infty \end{array} \right\rangle,$$

where $M = (m_{i,j})$ is the Coxeter matrix of Γ . Define the *Artin group* associated to Γ to be the group presented by

$$A = A(\Gamma) = \left\langle s_1, s_2, \dots, s_n \mid \begin{array}{l} \text{prod}_{m_{i,j}}(s_i, s_j) = \text{prod}_{m_{j,i}}(s_j, s_i) \\ \quad i \neq j \text{ and } m_{i,j} \neq \infty \end{array} \right\rangle.$$

If the group $W(\Gamma)$ is finite, then we say that $A(\Gamma)$ is an *Artin group of spherical type*. It is known that the Artin groups of spherical type have explicit solutions for the conjugacy problem (see [4], [10], [12], [13] and [21]). Moreover, we also know, for each of these groups, algorithms that compute a finite generating set for the centralizer of any element of $A(\Gamma)$ (see [11]). So Artin groups of spherical type are examples of groups in which we can apply our theory. A complete classification of these groups is given in [5]. Throughout this paper we will denote by $\mu : A(\Gamma) \rightarrow W(\Gamma)$ the canonical epimorphism.

An important example of a finite Coxeter group is the n th symmetric group, Σ_n . The group Σ_n is generated by the simple transpositions $\tau_i = (i, i+1)$, $i = 1, \dots, n-1$. The Coxeter relations in Σ_n are defined by the $(n-1) \times (n-1)$ Coxeter matrix $M = (m_{i,j})$, where $m_{i,i} = 1$ for all $i = 1, \dots, n-1$, $m_{i+1,i} = m_{i,i+1} = 3$ for all $i = 1, \dots, n-2$, and $m_{i,j} = 2$ for all $i, j = 1, \dots, n-1$, $|i-j| \geq 2$. The corresponding Coxeter graph is the graph A_{n-1} pictured in Figure 1 (see [5]).



FIGURE 1. Coxeter graph A_{n-1}

The Artin group associated to Σ_n is the braid group, $\mathcal{B}_n$, on n strands defined by Artin in [3]. Artin's presentation for the braid group $\mathcal{B}_n$ is

$$\mathcal{B}_n = \left\langle \sigma_1, \sigma_2, \dots, \sigma_{n-1} \mid \begin{array}{ll} \sigma_i \sigma_j = \sigma_j \sigma_i & (|i-j| \geq 2) \\ \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} & (i = 1, \dots, n-2) \end{array} \right\rangle.$$

The braid group $\mathcal{B}_n$ admits also a geometric interpretation as follows. Consider a disc D , and n distinct points $P_1, \dots, P_n$ in the interior of D . A *braid* b is the union of n disjoint paths $b_1, \dots, b_n$ in the cylinder $D \times [0, 1]$ such that, for all $i = 1, \dots, n$, the path b_i (the i th strand) goes monotonically in $t \in [0, 1]$ from $(P_i, 0)$ to some $(P_j, 1)$. We consider braids modulo isotopy, that is, two braids are equivalent if we can transform one into the other by deforming the strands, but keeping the ends fixed. Then the braid group is the group of isotopy classes of braids.

Note that the canonical epimorphism $\mu : \mathcal{B}_n \longrightarrow \Sigma_n$ is defined by: $\mu(\sigma_i) = \tau_i$. Let $b \in \mathcal{B}_n$. We say that b is a *pure braid* if $\mu(b) = 1$, or, equivalently, if the i th strand goes from $(P_i, 0)$ to $(P_i, 1)$, for all $i = 1, \dots, n$.

Example 1: Let $n \in \mathbb{N}$, $X \subset \{1, \dots, n\}$, and Σ_n be the n th symmetric group. Define $\Sigma_n(X)$ to be the subgroup of $\omega \in \Sigma_n$ such that, for all $i \in X$, $\omega(i) = i$. Define $\mathcal{B}_n(X) = \mu^{-1}(\Sigma_n(X))$. Briefly speaking, the group $\mathcal{B}_n(X)$ is the set of elements in $\mathcal{B}_n$ that are pure in the x th strand, for all $x \in X$.

Since the group Σ_n is finite, the triple $(\Sigma_n, \Sigma_n(X), \mu)$ verifies Condition **PC**, hence we can apply our theory to the subgroup $\mathcal{B}_n(X) < \mathcal{B}_n$, and shall obtain by this way a solution to the explicit conjugacy problem in $\mathcal{B}_n(X)$. Remark that $\mathcal{B}_n(X)$ is a finite index subgroup of $\mathcal{B}_n$, hence we can also give an algorithm to compute a finite generating set of the centralizer of any element of $\mathcal{B}_n(X)$. This example will be used later.

Example 2: The above construction has an easy extension to all spherical type Artin groups as follows. Pick a subgroup K' of W . Then, as W is finite, the triple (W, K', μ) verifies Condition **PC**, and we can again apply our theory to the subgroup $\mu^{-1}(K') < A$. A particular case is the following. Let W be a Coxeter group and let A be the corresponding Artin group. We define the *colored Artin group*, CA , as the kernel of the canonical epimorphism $\mu : A \longrightarrow W$. The colored Artin groups of spherical type appear in the theory of arrangements of hyperplanes as fundamental groups of complements of the so-called Coxeter arrangements (see [20]).

Example 3: The Artin group of type $\tilde{A}_{n-1}$ is the group associated to the Coxeter graph pictured in Figure 2, and the Artin group of type B_n




FIGURE 2. Coxeter graph $\tilde{A}_{n-1}$

has a presentation given by the Coxeter graph pictured in Figure 3. The Coxeter group $W(\tilde{A}_{n-1})$ is isomorphic to the infinite group $\mathbb{Z}^{n-1} \rtimes \Sigma_n$, so the Artin group $A(\tilde{A}_{n-1})$ is not of spherical type. Let $C_2 = \{\pm 1\}$ denote the

FIGURE 3. Coxeter graph B_n

cyclic group of order 2. On the other hand, the Coxeter group of type B_n is isomorphic to the semi-direct product $(C_2)^n \rtimes \Sigma_n$ which is finite, hence $A(B_n)$ is of spherical type. Let $\{a_1, \dots, a_n\}$ be the standard generators of $A(\tilde{A}_{n-1})$, and let $\{b_1, \dots, b_n\}$ be the standard generators of $A(B_n)$. Consider the homomorphism $\Phi : A(\tilde{A}_{n-1}) \longrightarrow A(B_n)$ defined by

$$\begin{aligned}\Phi(a_i) &= b_i, \quad i = 1, \dots, n-1 \\ \Phi(a_n) &= b_n^{-1} b_{n-1}^{-1} \cdots b_2^{-1} b_1 \cdots b_n.\end{aligned}$$

The homomorphism Φ is injective (see [1]) allowing us to see the group $A(\tilde{A}_n)$ as a subgroup of $A(B_n)$.

Define the homomorphism $\varphi : A(B_n) \longrightarrow \mathbb{Z}$ by $\varphi(b_n) = 1$ and $\varphi(b_i) = 0$ for $i < n$. In [16] it is proved that the group $A(\tilde{A}_n)$ is the inverse image of $\{1\}$ by φ . Now, because $\mathbb{Z}$ is abelian, hence it is endowed with a solution to the membership problem, and $\{1\}$ is finite, the triple $(\mathbb{Z}, \{1\}, \varphi)$ verifies Conditions **PC**. So, we can apply our theory to the pair $(A(\tilde{A}_{n-1}), A(B_n))$, and give an explicit solution to the conjugacy problem in $A(\tilde{A}_{n-1})$. Another solution to the conjugacy problem in $A(\tilde{A}_{n-1})$ can be derived from the facts that $A(\tilde{A}_{n-1})$ is biautomatic (see [7]), and that a biautomatic structure on a group provides a solution to the conjugacy problem (see [9]). The centralizers of some specific elements of $A(\tilde{A}_{n-1})$ are given in [8].

Example 4: Let $X = \{1, \dots, m\} \subset \{1, \dots, n\}$, and let $\pi_X : \mathcal{B}_n(X) \longrightarrow C\mathcal{B}_m$ be the homomorphism which sends a braid $b = (b_1, \dots, b_n) \in \mathcal{B}_n(X)$ to the pure braid $\pi_X(b) = (b_1, \dots, b_m)$. It is easily checked that π_X is a well-defined homomorphism. Define $\mathcal{IB}_n(X)$ as the kernel of the projection π_X . Note that $\mathcal{IB}_n(X)$ is the braid group on $n - m$ strands of the m punctured disc. Presentations for these groups are given in [17].

Assume $X = \{1, 2\}$. Then the Artin group of type $\tilde{C}_{n-1}$, defined by the Coxeter graph pictured in Figure 4, is isomorphic to $\mathcal{B}_n(X)$ (see [1]). By

FIGURE 4. Coxeter graph of $\tilde{C}_{n-1}$

Example 1, we have that the group $\mathcal{B}_n(X)$ has a solution to the explicit conjugacy problem, and we can compute a generating set of the centralizer of b for all $b \in \mathcal{B}_n(X)$. Moreover, the colored braid group $C\mathcal{B}_2$ is isomorphic to $\mathbb{Z}$, thus the triple $(C\mathcal{B}_2, \{1\}, \pi_X)$ verifies Conditions **PC**, hence we can

apply our theory to the subgroup $\mathcal{IB}_n(X) < \mathcal{B}_n(X)$, and obtain a solution to the explicit conjugacy problem in $A(\tilde{C}_{n-1})$. As far as I know, there is no other known solution to the conjugacy problem in $A(\tilde{C}_{n-1})$.

Beside the finite groups, another family of well-understood Coxeter groups are the so-called affine Coxeter groups which include, notably, the groups of type $\tilde{A}_n$ and $\tilde{C}_n$. The corresponding Artin groups, called *affine type Artin groups*, are poorly understood. Some other recent results concerning the groups $A(\tilde{A}_n)$ and $A(\tilde{C}_n)$ can be founded in [6], [7] and [8]. The previous techniques (Examples 1 and 3) give algorithms to treat the cases $\tilde{A}_n$ and $\tilde{C}_n$, but we do not know how to deal with the other affine type Artin groups.

Now, we show that it is not possible to apply our theory to the pair $(\mathcal{IB}_n(X), \mathcal{B}_n(X))$ for all $m \geq 5$. We do not know how to treat the cases $m = 3$ and $m = 4$.

Remark: If $m = n$ then $\mathcal{IB}_n(X)$ is the trivial group, hence we have a solution to the conjugacy problem. If $m = n - 1$ then $\mathcal{IB}_n(X) \cong F_m$ (the free group on m generators), thus we also have a solution to the conjugacy problem in $\mathcal{IB}_n(X)$. If $3 \leq m \leq n - 2$ then we do not know any solution to the conjugacy problem in $\mathcal{IB}_n(X)$.

Proposition 2.1. *Let $X = \{1, \dots, m\} \subset \{1, \dots, n\}$, $m \geq 5$. There is no triple (K, K', ϕ) such that K is a group endowed with a solution to the membership problem, K' is a subgroup of K , ϕ is a homomorphism $\phi : \mathcal{B}_n(X) \longrightarrow K$, and $\phi^{-1}(K') = \mathcal{IB}_n(X)$.*

Proof. Note that, if such a triple exists, then we can assume that the homomorphism ϕ is surjective. If ϕ is not surjective, and (K, K', ϕ) verifies Condition **PC**, then the triple $(\text{Im}(\phi), K', \phi)$ also verifies Condition **PC** and, in this case, ϕ is surjective.

Consider the natural embedding $\iota : \mathcal{B}_m \longrightarrow \mathcal{B}_n(X)$ defined by $\iota(\sigma_i) = \sigma_i$ for all $i = 1, \dots, m - 1$. Then ι restricts to a monomorphism $\iota : C\mathcal{B}_m \longrightarrow \mathcal{B}_n(X)$ which, actually, is a section of $\pi_X : \mathcal{B}_n(X) \longrightarrow C\mathcal{B}_m$. So, we have the split exact sequence,

$$1 \longrightarrow \mathcal{IB}_n(X) \hookrightarrow \mathcal{B}_n(X) \begin{array}{c} \xleftarrow{\iota} \\ \xrightarrow{\pi_X} \end{array} C\mathcal{B}_m \longrightarrow 1.$$

Define the homomorphism $\pi'_X : K \longrightarrow \mathcal{B}_m$ as follows. Let $v \in K$. Take $\tilde{v} \in \mathcal{B}_n(X)$ such that $\phi(\tilde{v}) = v$, and set $\pi'_X(v) = \pi_X(\tilde{v})$. Suppose that $\tilde{w}$ is another element of $\mathcal{B}_n(X)$ such that $\phi(\tilde{w}) = v$. Then $\phi(\tilde{w}\tilde{v}^{-1}) = 1$, thus, since $1 \in K'$, $\tilde{w}\tilde{v}^{-1} \in \phi^{-1}(K') = \mathcal{IB}_n(X)$, therefore $\pi_X(\tilde{w}\tilde{v}^{-1}) = 1$, that is, $\pi_X(\tilde{w}) = \pi_X(\tilde{v})$. This shows that the homomorphism π'_X is well-defined. Remark that, by construction, $\pi'_X \circ \phi = \pi_X$.

Now we prove that $\text{Ker}\pi'_X = K'$. Let $a \in K'$. Take $\tilde{a} \in \mathcal{IB}_n(X)$ such that $\phi(\tilde{a}) = a$. So, $\pi'_X(a) = \pi_X(\tilde{a}) = 1$, hence $a \in \text{Ker}\pi'_X$. Let $a \in \text{Ker}\pi'_X$. Then there exists $\tilde{a} \in \mathcal{B}_n(X)$ such that $\phi(\tilde{a}) = a$. Because $\pi_X(\tilde{a}) = \pi'_X(a) = 1$, we have $\tilde{a} \in \mathcal{IB}_n(X)$ and $\phi(\tilde{a}) = a \in K'$.

Let $\iota' = \phi \circ \iota$, and observe that ι' is a section of π'_X .

At this point, we have proved that the following diagram is commutative and that the lines are split exact sequences.

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \mathcal{IB}_n(X) & \hookrightarrow & \mathcal{B}_n(X) & \begin{array}{c} \xleftarrow{\iota} \\ \xrightarrow{\pi_X} \end{array} & \mathcal{CB}_m \longrightarrow 1 \\
 & & \downarrow \phi & & \downarrow \phi & & \downarrow id \\
 1 & \longrightarrow & K' & \hookrightarrow & K & \begin{array}{c} \xleftarrow{\iota'} \\ \xrightarrow{\pi'_X} \end{array} & \mathcal{CB}_m \longrightarrow 1
 \end{array}$$

Let F_2 be the free group on two generators, and let $R = F_2 \times F_2$. Theorem 1 of [18] tells that we can inject R in $\mathcal{CB}_m$, if $m \geq 5$, and Theorem 1 of [19] says that there is no solution to the membership problem in R . Now, $\mathcal{CB}_m$ can be embedded in K via ι' , thus R can be embedded in K , too, so, K has no solution to the membership problem. $\square$

3. THE CONJUGACY ALGORITHM

In this section we assume that G is a group, H is a subgroup of G , and (K, K', ϕ) is a triple that verifies Condition **PC**, and we prove an algorithm which solves the explicit conjugacy problem in H .

We start with a preliminary algorithm.

Algorithm 3.1. From conjugation in G to conjugation in H

Let $a, b \in H$, $h \in G$ such that $b = h^{-1}ah$. This algorithm will return FALSE, if a and b are not conjugated in H , and TRUE together with an element $h' \in H$ such that $b = (h')^{-1}ah'$, otherwise. This algorithm works as follows.

Let $a, b \in H$, and $h \in G$ such that $b = h^{-1}ah$. Recall that the group G is endowed with an algorithm that computes a finite generating set D of $C_G(a)$. By Condition **PC** the set $\phi(C_G(a)h) \cap K'$ is finite, and we can enumerate all its elements as follows. Observe that $\phi(C_G(a)) = \langle \phi(D) \rangle$. For each $k \in K'$, check whether $kh^{-1} \in \phi(C_G(a))$. This can be done because we know that $\phi(C_G(a))$ is generated by $\phi(D)$, and because K is endowed with a solution to the membership problem. Since K' is finite this process will stop. If $\phi(C_G(a)h) \cap K' = \emptyset$, then a and b are not conjugated in H . Indeed, if $b = (h')^{-1}ah'$ for some $h' \in H$, then $h'h^{-1} \in C_G(a)$, and $\phi(h') = \phi(h'h^{-1}h) \in K'$, so $\phi(C_G(a)h) \cap K' \neq \emptyset$. Suppose that $\phi(C_G(a)h) \cap K' \neq \emptyset$. Take an element $p \in \phi(C_G(a)h) \cap K'$ and write $p = \phi(d_1) \cdots \phi(d_q)\phi(h)$, where $d_1, \dots, d_q \in D$. Set $d = d_1 \cdots d_q$, and $h' = dh$. Then $\phi(h') = p \in K'$ (thus $h' \in H$), and $(h')^{-1}ah' = h^{-1}d^{-1}adh = h^{-1}ah = b$. We can resume this algorithm in the following way.

INPUT: A group G , a subgroup H of G , a triple (K, K', ϕ) that verifies Condition **PC**, two elements $a, b \in H$, an element $h \in G$ such that $b = h^{-1}ah$.

- (1) Compute a generating set D of $C_G(a)$.
- (2) For each $p \in K'$ do:
 - (a) If $p\phi(h^{-1}) \in \phi(C_G(a))$, then do:
 - (i) Write $p = \phi(d_1) \cdots \phi(d_q)\phi(h)$, where $d_1, \dots, d_q \in D$.
 - (ii) return TRUE and $h' = d_1 \cdots d_q h$.
- (3) return FALSE

OUTPUT: either FALSE if a and b are not conjugated in H , or TRUE together with an element $h' \in H$ such that $b = h'^{-1}ah'$, otherwise.

Algorithm 3.2. Explicit conjugacy algorithm in H

Given $a, b \in H \subset G$, apply the known conjugacy algorithm to a and b . If a and b are not conjugated as elements in G , then they cannot be conjugated in H . Otherwise, take $h \in G$ such that $h^{-1}ah = b$, and apply Algorithm 3.2.

INPUT: A group G , a subgroup H of G , a triple (K, K', ϕ) that verifies Condition **PC**, two elements $a, b \in H$.

- (1) For $a, b \in H$ test if they are conjugated in G . If they are not then return FALSE.
- (2) Let $h \in G$ be the output of the explicit conjugacy algorithm of G .
- (3) Apply Algorithm 3.1 to a, b and h and return the output of Algorithm 3.1

OUTPUT: either FALSE if a and b are not conjugated in H , or TRUE together with an element $h' \in H$ such that $b = h'^{-1}ah'$, otherwise.

4. CONSIDERATIONS ON THE COMPLEXITY OF THE ALGORITHMS

In general it is not possible to present an upper bound for the complexity of the algorithms presented in the previous section. There are several factors to take in charge. The first ones are the complexities of the conjugacy problem and of the algorithm to compute a generating set of the centralizers in G . These complexities, in the cases of the Artin groups of spherical type, are related to the size of a particular subset of the conjugacy class, the number of generators and the length of the word (see [10], [11] and [13]). Regarding the algorithms presented, we have also the following factors, the complexity of a solution to the membership problem in K , and the number of generators of the centralizers in G . Regarding this last factor we have some upper bounds in the case of the braid group (see [14]).

REFERENCES

- [1] D. Allcock, Braid pictures for artin groups, *Trans. A.M.S.* 354 (2002) 3455-3474.
- [2] I. Anshel, M. Anshel and D. Goldfeld, An algebraic method for public-key cryptography. *Math. Res. Lett.* **6**, No. 3-4 (1999), 287-291.
- [3] E. Artin, Theory of braids, *Ann. of Math.* (2) 48 (1947) 101-126
- [4] J. Birman, K. H. Ko and S. J. Lee, A new approach to the word and conjugacy problems in the braid groups, *Adv. Math.* **139**, No. 2 (1998), 322-353.
- [5] N. Bourbaki, Groupes et Algebres de Lie, Chapitres 4,5 et 6
- [6] R. Charney and J. Crisp, Automorphism groups of some affine and finite type Artin groups, Preprint arXiv:math.GR/0408412
- [7] R. Charney and D. Peifer, The $K(\pi, 1)$ conjecture for the affine braid groups, *Commentarii Math. Helv.* 78 (2003) 584-600
- [8] F. Digne, Presentations duales des groupes de tresses de type affine $\tilde{A}$, Preprint arXiv:math.GR/0402116
- [9] D. B. A. Epstein, J. W. Cannon, D. F. Holt, S. V. F. Levy, M. S. Paterson and W. P. Thurston, Braid Groups, Word processing in groups, Jones and Bartlett Publishers, Boston, MA, 1992.
- [10] N. Franco, J. Gonzalez-Meneses, Conjugacy problem for braid groups and Garside groups, *Journal of Algebra* 266 (1) 2003, 112-132.
- [11] N. Franco, J. Gonzalez-Meneses, Computation of centralizers in braid groups and Garside groups *Revista Matematica Iberoamericana*, 19 (2), 2003, 367-384.
- [12] F. A. Garside, The braid group and other groups, *Quart. J. Math. Oxford* **20** (1969), 235-154.
- [13] V. Gebhardt, A new aproach to the conjugacy problem in Garside groups, preprint.
- [14] J. Gonzalez-Meneses, B. Wiest, On the structure of the centralizer of a braid. To appear in the *Annales de l'ENS*.
- [15] K. H. Ko, S. J. Lee, J. H. Cheon, J. W. Han, J. Kang and C. Park, New public-key cryptosystem using braid groups. *Advances in cryptology—CRYPTO 2000* (Santa Barbara, CA), 166-183, *Lecture Notes in Comput. Sci.* **1880**, Springer, Berlin, 2000.
- [16] R. Kent and D. Peifer, A geometric and algebraic description of the annular braid groups, *Int. Jour. Alg. Comp.* 12 (2002), 85-97.
- [17] S. Lambropoulou, Braid structures in knot complements handlebodies and 3-manifolds, *Proceedings of the Conference "Knots in Hellas '98"*, Series of Knots and Everything, 24, World Scientific Press, 274-289 (2000).
- [18] T. A. Makanina, The occurrence problem for the braid group B_{n+1} when $n+1 \geq 5$. (Russian) *Mat. Zametki* 29 (1981), no. 1, 31-33, 154.
- [19] K. A. Mikhailova, The ocurrence problem for direct products of groups, (in Russian), *Mat. Sbornik*, v. 70 (1966), no. 2.
- [20] P. Orlik, H. Terao, Arrangements of Hyperplanes, *Grundlehren der Math. Wiss.* 300, Springer Verlag, 1992.
- [21] M. Picantin, The conjugacy problem in small Gaussian groups, *Comm. Algebra* **29**, No. 3 (2001), 1021-1039.

Nuno Franco,

R. Romão Ramalho 59, Departamento de Matemática, CIMA-UE, Universidade de Évora, 7000 Évora, Portugal

Institut de Mathématiques de Bourgogne, UMR 5584 du CNRS, Université de Bourgogne, B.P. 47870, 21078 Dijon cedex, France

E-mail: nmf@uevora.pt